

# CARDIOIDS AND TEICHMÜLLER SPACES

TOSHIYUKI SUGAWA

ABSTRACT. In this note, we give an expository account on the role played by cardioids in both Teichmüller spaces and in complex dynamics. Especially, we will observe that the shape of the Bers embedding of the Teichmüller space of a once-punctured rectangular torus converges to a cardioid in the sense of Carathéodory when the base surface is pinched along the meridian. A more general and complete result will be included in a forthcoming paper [21] of the author.

## 1. INTRODUCTION

A cardioid is a plane curve similar to  $\{e^{i\theta} - e^{2i\theta}/2 : \theta \in \mathbb{R}\}$ , see Figure 1. In this note, however, a *cardioid* means the (open) Jordan region bounded by a cardioid curve. We will call the *standard cardioid* the domain bounded by the above curve and denote it by  $C_0$ . Note that the function  $f_0(z) = z - z^2/2$  maps the unit disk univalently onto  $C_0$ . Conventionally, we denote by  $aC_0 + b$  the image of  $C_0$  under the similarity map  $z \mapsto az + b$ .

It is an accidental coincidence that the cardioid appears both in the Kleinian group theory and in complex dynamics. However, according to Sullivan's dictionary, the Bers embedding of the Teichmüller space of a Riemann surface (or a Fuchsian group) corresponds to the Mandelbrot set (or its generalization). Therefore, this coincidence might not be a surprising fact. As is well known, the Mandelbrot set contains a cardioid (the so-called main cardioid) as a connected component of its interior. On the other hand,




FIGURE 1. The cardioid

---

*Date:* August 30, 2006, *File:* teich\_HRI.tex.

*Key words and phrases.* cardioid, Teichmüller space, holomorphic motion, once-punctured torus.

The author was partially supported by the JSPS Grant-in-Aid for Scientific Research (B), 17340039.




FIGURE 2. The Mandelbrot set

as Kalme [8] pointed out, the cardioid appears in the universal Teichmüller space in a natural way.

In this note, we point out similarities between the cardioid in the Mandelbrot set and cardioids in the Bers embedded Teichmüller spaces. We emphasize on the role played by holomorphic motions to give a unified aspect on these matters. Moreover, we will observe that the shape of a one-dimensional Teichmüller space tends to that of a cardioid when the base Riemann surface goes to the boundary of the moduli space, at least in a special case. It appears that this is a phenomenon which has not been observed in the literature.

The author would like to thank the organizers of the International Workshop on Teichmüller Theory and Moduli Problems held at the Harish-Chandra Research Institute, Allahabad, India, for giving him a chance to consider this matter seriously. Without their kind invitation to the workshop, this work would not have been carried out due to the author's busyness and laziness.

## 2. CARDIOID IN MANDELBROT SET

Even polynomials of the simple form  $P_c(z) = z^2 + c$  are not fully understood in the context of iterations, that is, in the theory of complex dynamics. We denote by  $P_c^n$  the  $n$ -th iterate of  $P_c$ , namely,  $P_c^n = P_c^{n-1} \circ P_c$  for  $n = 2, 3, \dots$ . Here  $c$  is a complex parameter. The boundary of the set  $K_c = \{z \in \mathbb{C} : \{P_c^n(z)\}_{n=1,2,\dots} \text{ is bounded}\}$  is called the Julia set and denoted by  $J_c$ . The Mandelbrot set  $M$  is defined to be the set of  $c \in \mathbb{C}$  for which  $J_c$  is connected (see Figure 2). It is known that  $J_c$  is either connected or a Cantor set. Also, it should be noted that  $c \notin M$  iff  $0 \in K_c$ .

An *attracting fixed point* of  $P_c$  is a point  $\alpha \in \mathbb{C}$  such that  $P_c(\alpha) = \alpha$  and  $|P_c'(\alpha)| < 1$ . It is easily seen that a neighbourhood of an attracting fixed point of  $P_c$  is contained in  $K_c$ . In particular,  $c \in M$ . A fixed point  $\alpha$  of  $P_c$  must satisfy the relation  $P_c(\alpha) - \alpha = \alpha^2 - \alpha + c = 0$ . If we set  $\lambda = P_c'(\alpha) = 2\alpha$ , then  $c = \lambda/2 - (\lambda/2)^2$ . We now conclude that  $P_c$  has an

attracting fixed point iff  $c \in D = \{\lambda/2 - (\lambda/2)^2 : \lambda \in \mathbb{D}\}$ . Here and hereafter,  $\mathbb{D}$  denotes the unit disk  $\{z \in \mathbb{C} : |z| < 1\}$ . Therefore,  $D$  is contained in the Mandelbrot set. Indeed, it is easy to see that  $D$  is a component of the interior of  $M$  and  $D = (1/2)C_0$ . The component  $D$  is called the main cardioid of  $M$ .

We expect that the Julia set  $J_c$  varies holomorphically (in a sense) on the parameter  $c \in D$ . Indeed, a repelling periodic point  $z_c$  of  $P_c$  is a holomorphic function of  $c \in D$  (if one takes a suitable branch). In this way, we obtain a function  $f_c(z_0) = z_c$  on  $R_0 = \{e^{2\pi i\theta} : 2^n\theta \in \mathbb{Z} \exists n \in \mathbb{N}\}$  for each  $c \in D$ . Since these repelling periodic points do not collide for  $c \in D$ , the function  $f_c$  is injective on  $R_0$ . Thus the map  $(c, z) \mapsto f_c(z)$  gives a holomorphic motion of  $R_0$  over  $D$ . Since  $R_0$  is dense in  $\partial\mathbb{D}$ , the holomorphic motion of  $R_0$  extends to that of  $\partial\mathbb{D}$  (see [11]), which will be still denoted by the same letter  $f_c(z)$ . Since the set of repelling periodic points is dense in the Julia set, one has the relation  $J_c = f_c(\partial\mathbb{D})$  for  $c \in D$ . By a theorem of Mañé-Sad-Sullivan [11], the map  $f_c$  can be extended to a quasiconformal map of  $\widehat{\mathbb{C}}$  and thus  $J_c = f_c(\partial\mathbb{D})$  is known to be a quasidisk for  $c \in D$ . For details and interesting figures, see Astala and Martin [2].

It is known that near the cusp point  $1/4$  of the main cardioid  $D$  rather complicated phenomena of phase transition are observed, which are called the parabolic explosion. See [3] for details and references.

### 3. BERS EMBEDDING OF TEICHMÜLLER SPACE

We give minimal basics in the theory of Teichmüller spaces to state and to prove our result. For details, see [14] or [7].

In the following, we denote by  $\mathbb{H}$  the upper half-plane  $\{z \in \mathbb{C} : \text{Im } z > 0\}$  and by  $\mathbb{H}^*$  the lower half-plane  $\{z \in \mathbb{C} : \text{Im } z < 0\}$ . Let  $\Gamma$  be a Fuchsian group acting on  $\mathbb{H}$ , namely,  $\Gamma$  is a discrete subgroup of  $\text{PSL}(2, \mathbb{R})$ . We denote by  $\text{Belt}(\mathbb{H}, \Gamma)$  the space of Beltrami coefficient for  $\Gamma$  on  $\mathbb{H}$ , more precisely,

$$\text{Belt}(\mathbb{H}, \Gamma) = \{\mu \in L^\infty(\mathbb{H}) : \|\mu\|_\infty < 1, (\mu \circ \gamma) \frac{\overline{\gamma'}}{\gamma'} = \mu \ \forall \gamma \in \Gamma\}.$$

For  $\mu \in \text{Belt}(\mathbb{H}, \Gamma)$ , we denote by  $f^\mu$  the quasiconformal map  $f$  of  $\widehat{\mathbb{C}} = \mathbb{C} \cup \{\infty\}$  which is determined by the Beltrami equations  $f_{\bar{z}} = \mu f_z$  in  $\mathbb{H}$  and  $f_{\bar{z}} = 0$  in  $\mathbb{H}^*$  and normalization conditions  $f(0) = 0, f(1) = 1$  and  $f(\infty) = \infty$ . Note that  $f^\mu \Gamma (f^\mu)^{-1}$  is a Kleinian group acting on  $f^\mu(\mathbb{H})$  and  $f^\mu(\mathbb{H}^*)$  properly discontinuously.

Noting that  $f^\mu$  is conformal on  $\mathbb{H}^*$ , we define a holomorphic function  $\Phi(\mu)$  on  $\mathbb{H}^*$  by  $\Phi(\mu) = S_{f^\mu|_{\mathbb{H}^*}}$ , where  $S_f$  stands for the Schwarzian derivative of  $f$ :

$$S_f = \left(\frac{f''}{f'}\right)' - \frac{1}{2} \left(\frac{f'}{f}\right)^2 = \frac{f'''}{f'} - \frac{3}{2} \left(\frac{f'}{f}\right)^2.$$

It is convenient to measure the Schwarzian derivative by the norm

$$\|\varphi\|_{\mathbb{H}^*} = \sup_{z \in \mathbb{H}^*} (-2 \text{Im } z)^2 |\varphi(z)|.$$

We denote by  $B_2(\mathbb{H}^*)$  the Banach space consisting of analytic functions  $\varphi$  on  $\mathbb{H}^*$  with  $\|\varphi\|_{\mathbb{H}^*} < \infty$ . Let

$$B_2(\mathbb{H}^*, \Gamma) = \{\varphi \in B_2(\mathbb{H}^*) : (\varphi \circ \gamma)(\gamma')^2 = \varphi \ \forall \gamma \in \Gamma\},$$

which is a closed subspace of  $B_2(\mathbb{H}^*)$ .

It is known that  $\Phi(\mu) \in B_2(\mathbb{H}^*, \Gamma)$  for  $\mu \in \text{Belt}(\mathbb{H}, \Gamma)$  and that the image  $\Phi(\text{Belt}(\mathbb{H}, \Gamma))$  coincides with the Bers embedding of the Teichmüller space of  $\Gamma$ . We set  $\text{Teich}(\Gamma) = \Phi(\text{Belt}(\mathbb{H}, \Gamma))$  and we identify it with the Teichmüller space of  $\Gamma$  (or equivalently, of the orbifold  $\mathbb{H}/\Gamma$ ). Note that  $\text{Teich}(\Gamma)$  is a bounded domain in  $B_2(\mathbb{H}^*, \Gamma)$ . The map  $\Phi_\Gamma = \Phi : \text{Belt}(\mathbb{H}, \Gamma) \rightarrow \text{Teich}(\Gamma)$  is called the Bers projection and known to be a holomorphic submersion. In the case when  $\Gamma$  is the trivial group 1, the set  $\text{Teich}(1)$  is called the *universal Teichmüller space*.

We denote by  $o(\Gamma)$  and  $i(\Gamma)$  the outer and inner radii of the Teichmüller space  $\text{Teich}(\Gamma)$ , in other words,  $o(\Gamma)$  is the smallest number  $r$  with  $\text{Teich}(\Gamma) \subset \{\varphi \in B_2(\mathbb{H}^*, \Gamma) : \|\varphi\|_{\mathbb{H}^*} < r\}$  and  $i(\Gamma)$  is the largest number  $r$  with  $\text{Teich}(\Gamma) \supset \{\varphi \in B_2(\mathbb{H}^*, \Gamma) : \|\varphi\|_{\mathbb{H}^*} < r\}$ .

It is well known that  $i(1) = 2$  and  $o(1) = 6$  for trivial group 1, and hence,  $2 \leq i(\Gamma) \leq o(\Gamma) \leq 6$  for an arbitrary Fuchsian group  $\Gamma$  unless  $\text{Teich}(\Gamma)$  is a singleton. For a cofinite Fuchsian group  $\Gamma$ , it is also known that  $i(\Gamma) > 2$  (cf. [16]) and  $o(\Gamma) < 6$  (due to Sekigawa [20]). Furthermore, Nakanishi and Yamamoto [18] and Nakanishi and Velling [17] gave characterizing conditions for  $\Gamma$  to satisfy  $o(\Gamma) = 6$  and  $i(\Gamma) = 2$ , respectively. It is remarkable that their conditions are coincident. In particular,  $o(\Gamma) = 6$  iff  $i(\Gamma) = 2$ .

#### 4. KALME'S OBSERVATION

Kalme [8] made an interesting observation on a special holomorphic family of quasiconformal maps of the Riemann sphere. Following [8], we present some facts related to our investigation.

For a complex number  $\alpha \in \mathbb{C}$ , consider the function

$$F_\alpha(z) = z^\alpha = e^{\alpha \log z}$$

on  $\mathbb{H}^*$ , where the branch of  $\log z$  is taken so that  $-\pi < \arg z = \text{Im} \log z < 0$  for  $z \in \mathbb{H}^*$ . Since  $F'_\alpha(z) = \alpha z^{\alpha-1} \neq 0$ , the map  $F_\alpha$  is locally univalent for  $\alpha \neq 0$ . A characterization of univalence of  $F_\alpha$  is known.

**Lemma 1** (Royster [19]). *Let  $\alpha \in \mathbb{C} \setminus \{0\}$ . Then,  $F_\alpha$  is univalent in  $\mathbb{H}^*$  iff  $|\alpha - 1| \leq 1$  or  $|\alpha + 1| \leq 1$ .*

An easy computation gives

$$S_{F_\alpha}(z) = \frac{1 - \alpha^2}{2z^2} = 2(1 - \alpha^2)\varphi_0^*(z),$$

where  $\varphi_0^* \in B_2(\mathbb{H}^*)$  is given by  $\varphi_0^*(z) = 1/(4z^2)$ . Note that  $\|\varphi_0^*\|_{\mathbb{H}^*} = 1$ .

An interesting fact is that the holomorphic motion  $f_t = F_{1-t}$  ( $|t| < 1$ ) of  $\mathbb{H}^*$  can be extended to that of the Riemann sphere in an explicit way:

$$f_t(z) = \begin{cases} z\bar{z}^{-t} & z \in \overline{\mathbb{H}} \\ z^{1-t} & z \in \mathbb{H}^*. \end{cases}$$

Note that  $f_t$  is normalized and has the Beltrami coefficient

$$\mu_{f_t}(z) = -t \frac{z}{\bar{z}} = -t \frac{|\varphi_0(z)|}{\varphi_0(z)}$$

for  $z \in \mathbb{H}$ , where  $\varphi_0(z) = \overline{\varphi_0^*(\bar{z})} = 1/(4z^2)$  for  $z \in \mathbb{H}$ . This is of the form of a Teichmüller differential on  $\mathbb{H}$  and, by definition of  $\Phi$ , one has the relation

$$(4.1) \quad \Phi(-t\mu_0) = 2(1 - (1 - t)^2)\varphi_0^* = g_0(t)\varphi_0^*,$$

where

$$\mu_0 = \frac{|\varphi_0|}{\varphi_0} \quad \text{and} \quad g_0(t) = 4 \left( t - \frac{t^2}{2} \right).$$

We can now see that the intersection of  $\text{Teich}(1)$  with the linear span of  $\varphi_0^*$  is precisely the cardioid  $4C_0$  (times  $\varphi_0^*$ ).

Hille [6] observed that for  $\varepsilon \in \mathbb{R} \setminus \{0\}$ ,  $F_{i\varepsilon}$  is a universal covering projection onto an annulus, and hence, is never univalent whereas  $S_{F_{i\varepsilon}}(z) = (1 + \varepsilon^2)/(2z^2)$  and thus the norm  $\|S_{F_{i\varepsilon}}\|_{\mathbb{H}^*}$  tends to 2 as  $\varepsilon \rightarrow 0$ . We point out that the point  $S_{F_{i\varepsilon}}$  approaches to the inward cusp of the above cardioid  $4C_0\varphi_0^*$  from the outside.

## 5. ONE-DIMENSIONAL TEICHMÜLLER SPACES.

From now on, we restrict ourselves on one-dimensional Teichmüller spaces. For simplicity, we assume Fuchsian groups to be torsion-free. Then  $\dim \text{Teich}(\Gamma) = 1$  iff the signature of  $\Gamma$  is  $(0,4)$  or  $(1,1)$ . Since to each Fuchsian group  $\Gamma$  of signature  $(0,4)$ , there corresponds a Fuchsian group  $\Gamma'$  of signature  $(1,1)$  such that  $\text{Teich}(\Gamma) = \text{Teich}(\Gamma')$  (cf. [9]), without loss of generality we may further assume that  $\Gamma$  is of signature  $(1,1)$ . Then the quotient Riemann surface  $\mathbb{H}/\Gamma$  is represented by the once-punctured torus of the form  $T = (\mathbb{C} \setminus \Omega)/\Omega$ , where

$$\Omega = \langle \omega_1, \omega_2 \rangle_{\mathbb{Z}} = \{m\omega_1 + n\omega_2 : m, n \in \mathbb{Z}\}$$

and  $\omega_1, \omega_2 \in \mathbb{C}$  with  $\text{Im}(\omega_2/\omega_1) > 0$ . We also write  $\text{Teich}(T)$  for  $\text{Teich}(\Gamma)$  when  $T = \mathbb{H}/\Gamma$ .

In recent years, much progress has been made in the study of one-dimensional Teichmüller spaces. For instance, Minsky [12] proved that the Bers embedding of a one-dimensional Teichmüller space is a Jordan domain. By using Minsky's method, Miyachi [13] showed that the one-dimensional Teichmüller space is "cusp-shaped" at every boundary point corresponding to a cusp. Computer graphics of the Bers embeddings are presented by [9] and [10]. Also, the inner and outer radii of the Teichmüller space of a once-punctured square torus are numerically computed in [22].

In [21], the following result is proved.

**Theorem 2.** *The Bers embedding  $\text{Teich}(T)$  of the Teichmüller space of a once-punctured torus  $T$  converges to a cardioid in the sense of Carathéodory (under suitable identification of  $B_2(T) = B_2(\mathbb{D}, \Gamma)$  with  $\mathbb{C}$ ), when  $T$  goes to infinity in the moduli space*

There is some technicality with the suitable choice of the basis of the vector space  $B_2(T)$  so that one can identify  $B_2(T)$  with  $\mathbb{C}$ . Therefore, in this note, we restrict ourselves to the case when  $T$  is a rectangular torus for the sake of simplicity.

Let  $\Omega_\lambda$  be the lattice generated by  $\lambda > 0$  and  $i = \sqrt{-1}$  over  $\mathbb{Z}$ , namely,  $\Omega_\lambda = \{m + ni : m, n \in \mathbb{Z}\}$ . We set  $T_\lambda = (\mathbb{C} \setminus \Omega_\lambda)/\Omega_\lambda$ . We denote by  $\pi_\lambda(\zeta) = [\zeta]$  the canonical projection  $\mathbb{C} \setminus \Omega_\lambda \rightarrow T_\lambda$ .

Take a holomorphic universal covering projection  $p_\lambda$  of the upper half-plane  $\mathbb{H}$  onto  $\mathbb{C} \setminus \Omega_\lambda$  so that  $p_\lambda(i) = a_\lambda$  and  $p'_\lambda(i)/i > 0$ . Then  $q_\lambda = \pi_\lambda \circ p_\lambda$  is a holomorphic universal

covering projection of  $\mathbb{H}$  onto  $T_\lambda$ . Let  $\Gamma_\lambda$  be the covering transformation group of  $q_\lambda : \mathbb{H} \rightarrow T_\lambda$ . Note that  $\mathbb{H}/\Gamma_\lambda = T_\lambda$ .

Let  $\tilde{\varphi}_\lambda$  be the pullback of the quadratic differential  $d\zeta^2$  on  $\mathbb{C} \setminus \Omega_\lambda$  under the map  $p_\lambda$ , namely,  $\tilde{\varphi}_\lambda = (p'_\lambda)^2$ . Further let

$$\varphi_\lambda = (\|\tilde{\varphi}_\lambda\|_{\mathbb{H}})^{-1} \tilde{\varphi}_\lambda.$$

Then, by definition,  $\varphi_\lambda \in B_2(\mathbb{H}, \Gamma_\lambda)$  and  $\|\varphi_\lambda\|_{\mathbb{H}} = 1$ .

Set also

$$\varphi_\lambda^*(z) = \overline{\varphi_\lambda(\bar{z})}.$$

Then  $\varphi_\lambda^* \in B_2(\mathbb{H}^*, \Gamma_\lambda)$  and  $\|\varphi_\lambda^*\|_{\mathbb{H}^*} = 1$ .

Since  $d\zeta^2$  on  $\mathbb{C} \setminus \Omega_\lambda$  projects to a nontrivial holomorphic quadratic differential on  $T_\lambda$  via  $\pi_\lambda$ , the vector space  $B_2(\mathbb{H}^*, \Gamma_\lambda)$  is spanned by  $\varphi_\lambda^*$ . Let

$$U_\lambda = \{w \in \mathbb{C} : w\varphi_\lambda^* \in \text{Teich}(\Gamma_\lambda)\}.$$

Then, our result can be stated as in the following.

**Theorem 3.** *The domain  $U_\lambda$  converges to the cardioid  $4C_0$  in the sense of Carathéodory as  $\lambda \rightarrow 0+$ .*

Since  $U_\lambda$  is a simply connected bounded domain (indeed, a Jordan domain by Minsky's theorem), there exists a conformal homeomorphism  $g_\lambda$  of the unit disk  $\mathbb{D}$  onto  $U_\lambda$  such that  $g_\lambda(0) = 0$  and  $g'_\lambda(0) > 0$ . Recall that  $g_0(t) = 2(1 - (t - 1)^2)$  is a conformal map of  $\mathbb{D}$  onto the cardioid  $4C_0$ . Then, the last theorem can be restated in the following way.

**Theorem 4.** *The conformal map  $g_\lambda$  converges to the map  $g_0$  locally uniformly on  $\mathbb{D}$ .*

We illustrate the shape of the boundary of  $\text{Teich}(\Gamma_\lambda)$  for several values of  $\lambda$ . We remark that the boundary curve is drawn very roughly near the “main” cusp when  $\lambda$  is small. This is simply because of our algorithm of computation.

## 6. PROOF OF THEOREM 4

Let  $\rho_{\mathbb{C} \setminus \Omega_\lambda}(\zeta)|d\zeta|$  and  $\rho_{T_\lambda}([\zeta])|d\zeta|$  be the hyperbolic metrics on  $\mathbb{C} \setminus \Omega_\lambda$  and  $T_\lambda$ , respectively. The density functions (the hyperbolic densities)  $\rho_{\mathbb{C} \setminus \Omega_\lambda}(\zeta)|d\zeta|$  and  $\rho_{T_\lambda}([\zeta])$  are characterized by the relations  $\rho_{T_\lambda}([\zeta]) = \rho_{\mathbb{C} \setminus \Omega_\lambda}(\zeta)$  and  $\rho_{\mathbb{C} \setminus \Omega_\lambda}(p_\lambda(z))|p'_\lambda(z)| = \rho_{\mathbb{H}}(z) = 1/(2\text{Im } z)$ . The following assertion can be found (in a more general form) in [22].








FIGURE 3. Bers embedding for the punctured square torus ( $\lambda = 1$ )

FIGURE 4. Bers embedding for small  $\lambda$ FIGURE 5. Bers embedding for very small  $\lambda$ FIGURE 6. Bers embedding for very very small  $\lambda$ FIGURE 7. The cardioid  $4C_0 = \{2(1 - \alpha^2) : |\alpha - 1| < 1\}$ .

**Lemma 5.** *The hyperbolic density  $\rho_{T_\lambda}([\zeta])$  of  $T_\lambda$  takes its minimum at the point  $[a_\lambda] = [(\lambda + i)/2]$ .*

**Corollary 6.** *The supremum of  $(2\operatorname{Im} z)^2 |\varphi_\lambda(z)|$  over  $z \in \mathbb{H}$  is attained at  $z = i$ .*

*Proof.* Letting  $\zeta = p_\lambda(z)$ , we have the relation

$$(2\operatorname{Im} z)^2 |\varphi_\lambda(z)| = \rho_{\mathbb{C} \setminus \Omega_\lambda}(\zeta)^{-2},$$

which is known to take its maximum at  $\zeta = a_\lambda = p_\lambda(i)$  by the last lemma.  $\square$

By the normalization of  $\varphi_\lambda$ , we have thus the relation  $4|\varphi_\lambda(i)| = \|\varphi_\lambda\|_{\mathbb{H}} = 1$ . On the other hand,  $\tilde{\varphi}_\lambda(i) = p'_\lambda(i)^2 < 0$ . Therefore,  $\varphi_\lambda(i) = \tilde{\varphi}_\lambda(i)/\|\tilde{\varphi}_\lambda\|_{\mathbb{H}} < 0$  and thus

$$(6.1) \quad \varphi_\lambda(i) = -\frac{1}{4}.$$

We also note the following fact, which will be used later.

**Lemma 7.** *The hyperbolic density  $\rho_{\mathbb{C} \setminus \Omega_\lambda}(\xi + i/2)$  converges to  $\pi/2$  uniformly in  $\xi \in \mathbb{R}$  as  $\lambda \rightarrow 0+$ .*

*Proof.* Observe that  $\mathbb{C} \setminus \Omega_\lambda$  converges to the parallel strip  $S = \{\zeta : 0 < \operatorname{Im} \zeta < 1\}$  with respect to the point  $i/2$  in the sense of Carathéodory when  $\lambda \rightarrow 0+$ . Thus, the hyperbolic density  $\rho_{\mathbb{C} \setminus \Omega_\lambda}$  converges to  $\rho_S$  locally uniformly on  $S$  (cf. [5]). Since  $\rho_{\mathbb{C} \setminus \Omega_\lambda}(\xi + i/2)$  is a periodic function of  $\xi$  with period  $\lambda$  and since  $\rho_S(\xi + i/2) = \pi/2$ , the assertion follows.  $\square$

We now define the curves

$$\begin{aligned} \alpha_\lambda(s) &= [a_\lambda + \lambda s] \quad (0 \leq s \leq 1) \\ \beta_\lambda(s) &= [a_\lambda + is] \quad (0 \leq s \leq 1) \end{aligned}$$

on  $T_\lambda$ , where we recall that  $a_\lambda = (\lambda + i)/2$ . Then, by the obvious symmetry of  $T_\lambda$ , these are simple hyperbolic geodesics of  $T_\lambda$  (see Figure 8).

We take the lifts  $\hat{\alpha}_\lambda$  and  $\hat{\beta}_\lambda$  of them starting at  $i$  via the covering projection  $q_\lambda : \mathbb{H} \rightarrow T_\lambda$ . We denote by  $A_\lambda$  and  $B_\lambda$  the elements of  $\Gamma_\lambda \subset \operatorname{PSL}(2, \mathbb{R})$  corresponding to  $\hat{\alpha}_\lambda$  and  $\hat{\beta}_\lambda$ , respectively. In other words,  $\hat{\alpha}_\lambda(1) = A_\lambda(\hat{\alpha}_\lambda(0)) = A_\lambda(i)$  and  $\hat{\beta}_\lambda(1) = B_\lambda(\hat{\beta}_\lambda(0)) = B_\lambda(i)$ . See Figure 9.

Let  $\sigma$  and  $\tau$  be the hyperbolic lengths of curves  $\alpha_\lambda$  and  $\beta_\lambda$ , respectively. Then  $A_\lambda$  and  $B_\lambda$  can be represented by

$$A_\lambda = \pm \begin{pmatrix} e^\sigma & 0 \\ 0 & e^{-\sigma} \end{pmatrix} \quad \text{and} \quad B_\lambda = \pm \begin{pmatrix} \cosh \tau & \sinh \tau \\ \sinh \tau & \cosh \tau \end{pmatrix}.$$

Since the commutator  $[A_\lambda, B_\lambda]$  must be parabolic, the following relation is required:

$$\sinh \sigma \sinh \tau = 1.$$

We remark that this sort of relation was already found by Hayman [4, §7].

We note the following fact.

**Lemma 8.** *The hyperbolic length  $\sigma$  of  $\alpha_\lambda$  tends to 0 as  $\lambda \rightarrow 0+$ .*






FIGURE 8. Geodesics on  $T_\lambda$




FIGURE 9. Lifts of curves on  $\mathbb{H}$

*Proof.* By definition,  $\sigma$  can be computed by

$$\sigma = \int_0^\lambda \rho_{T_\lambda}([\xi + i/2])d\xi = \int_0^\lambda \rho_{\mathbb{C} \setminus \Omega_\lambda}(\xi + i/2)d\xi.$$

By Lemma 7, we see that  $\sigma \sim \pi\lambda/2$  as  $\lambda \rightarrow 0 +$ . In particular, the assertion follows.  $\square$

A crucial result is the following variant of a proposition due to T. Nakanishi [15, Proposition 3.1].

**Lemma 9** (Nakanishi's lemma). *Let  $G_n$  be a sequence of Fuchsian groups acting on  $\mathbb{H}$  each of which contains a hyperbolic element of the form  $z \mapsto M_n z$  such that  $M_n \rightarrow 1$  as  $n \rightarrow \infty$ . Further let  $\varphi_n$  be an element of  $B_2(\mathbb{H}, G_n)$  such that  $\varphi_n$  converges locally uniformly to a holomorphic function  $\varphi_\infty$  on  $\mathbb{H}$ . Then  $\varphi_\infty(z) = c/z^2$  for some constant  $c$ .*

*Proof.* Write  $\varphi_n(z) = P_n(z)/z^2$  and  $\varphi_\infty(z) = P_\infty(z)/z^2$ . Then  $P_n \rightarrow P_\infty$  locally uniformly. Since  $\varphi_n(M_n z)M_n^2 = \varphi_n(z)$ , we have  $P_n(M_n^k z) = P_n(z)$  for any  $k \in \mathbb{Z}$ . Letting  $n \rightarrow \infty$ , we can see that  $P_\infty$  is constant along the positive imaginary axis, which implies that  $P_\infty$  is constant.  $\square$

**Lemma 10.** *The quadratic differential  $\varphi_\lambda(z)$  converges to  $\varphi_0(z) = 1/(4z^2)$  locally uniformly on  $\mathbb{H}$  as  $\lambda \rightarrow 0+$ .*

*Proof.* Since  $\|\varphi_\lambda\|_{\mathbb{H}} = 1$ , the family  $\{\varphi_\lambda\}$  is locally bounded on  $\mathbb{H}$  and thus normal. Let  $\lambda_n$  be any sequence of positive numbers tending to 0 such that  $\varphi_{\lambda_n}$  converges locally uniformly on  $\mathbb{H}$ . Then Lemmas 8 and 9 imply that the limit function has the form  $c/z^2$ . On the other hand, by (6.1), the constant  $c$  must be  $1/4$ , which is independent of the sequence  $\lambda_n$ . Thus a standard argument gives us the local uniform convergence of  $\varphi_\lambda$  to  $1/(4z^2)$ .  $\square$

Let

$$\mu_\lambda = \frac{|\varphi_\lambda|}{\varphi_\lambda}.$$

Then  $t\mu_\lambda \in \text{Belt}(\mathbb{H}, \Gamma_\lambda)$  for every  $t \in \mathbb{D}$ . Since  $\Phi(-t\mu_\lambda) \in B_2(\mathbb{H}^*, \Gamma_\lambda)$ , we can write  $\Phi(-t\mu_\lambda) = c\varphi_\lambda^*$  for some  $c \in \mathbb{C}$  for each  $t \in \mathbb{D}$ . We define the function  $h_\lambda : \mathbb{D} \rightarrow \mathbb{C}$  by the relation

$$\Phi(-t\mu_\lambda) = h_\lambda(t)\varphi_\lambda^*.$$

We now show the following.

**Lemma 11.** *The function  $h_\lambda$  is a conformal homeomorphism of  $\mathbb{D}$  onto  $U_\lambda$  satisfying  $h_\lambda(0) = 0$ ,  $\overline{h_\lambda(t)} = h_\lambda(\bar{t})$ , and, in particular,  $h'_\lambda(0) \in \mathbb{R} \setminus \{0\}$ .*

*Proof.* Since the map  $t \mapsto \Phi_{\Gamma_\lambda}(-t\mu_\lambda)$  is of the form of a Teichmüller map,  $h_\lambda$  is known to be a proper holomorphic injection of  $\mathbb{D}$  into  $U_\lambda$  (see, for instance, [14]). Since  $\dim \text{Teich}(\Gamma_\lambda) = 1$ , this map is indeed a conformal homeomorphism.

By the symmetry of  $\Omega_\lambda$ , we have the relation  $\overline{p_\lambda(-\bar{z})} - a_\lambda + a_\lambda = p_\lambda(z)$ . Therefore,  $\varphi_\lambda = (p'_\lambda)^2$  satisfies  $\overline{\varphi_\lambda(-\bar{z})} = \varphi_\lambda(z)$ . Hence,  $\overline{\mu_\lambda(-\bar{z})} = \mu_\lambda(z)$ . This implies  $\overline{\Phi(\bar{t}\mu_\lambda)(-\bar{z})} = \Phi(t\mu_\lambda)(z)$ , which is equivalent to  $\overline{h_\lambda(\bar{t})\varphi_\lambda^*(-\bar{z})} = h_\lambda(t)\varphi_\lambda^*(z)$ . Since  $\overline{\varphi_\lambda^*(-\bar{z})} = \varphi_\lambda^*(z)$ , we have the relation  $\overline{h_\lambda(\bar{t})} = h_\lambda(t)$ .  $\square$

*Remark.* The quasiconformal deformation of  $T_\lambda$  corresponding to  $\Phi(-t\mu_\lambda)$  can be constructed explicitly. This holomorphic motion of  $\mathbb{H}^*$  is induced by the holomorphic motion

of  $\mathbb{C} \setminus \Omega_\lambda$  given by

$$\mathbb{D} \times (\mathbb{C} \setminus \Omega_\lambda) \ni (t, \zeta) \mapsto \frac{\zeta - t\bar{\zeta}}{1+t}.$$

See [21] for the details.

By Lemma 10, we see that the Teichmüller differential  $\mu_\lambda = |\varphi_\lambda|/\varphi_\lambda$  converges pointwise to  $\mu_0 = |\varphi_0|/\varphi_0 = z/\bar{z}$ . We now use the following basic property of quasiconformal maps. The proof of it can be read from a paper of Ahlfors and Bers [1]. See also [21] for a detailed account.

**Proposition 12.** *Let  $\mu_n$  be a sequence of measurable functions on  $\mathbb{C}$  such that  $\|\mu_n\|_\infty \leq 1$  and  $\mu_n \rightarrow \mu$  a.e. Then the normalized  $t\mu_n$ -quasiconformal map  $f^{t\mu_n}(z)$  converges to  $f^{t\mu}(z)$  locally uniformly in  $(t, z) \in \mathbb{D} \times \mathbb{C}$ .*

By the above proposition,  $f^{-t\mu_\lambda}(z)$  converges to  $f^{-t\mu_0}(z)$  locally uniformly in  $(t, z) \in \mathbb{D} \times \mathbb{C}$  as  $\lambda \rightarrow 0+$ . Since  $f^{-t\mu_\lambda}$  is analytic in  $\mathbb{H}^*$ , the Weierstrass double series theorem implies that

$$\Phi(-t\mu_\lambda)(z) = S_{f^{-t\mu_\lambda}}(z) \rightarrow S_{f^{-t\mu_0}}(z) = \Phi(-t\mu_0)(z)$$

locally uniformly in  $(t, z) \in \mathbb{D} \times \mathbb{H}^*$ . By definition of  $h_\lambda$  and (4.1), the above convergence is equivalent to the locally uniform convergence

$$h_\lambda(t)\varphi_\lambda^*(z) \rightarrow g_0(t)\varphi_0^*(z).$$

Since  $\varphi_\lambda^* \rightarrow \varphi_0^*$  by Lemma 10, we now see that  $h_\lambda \rightarrow g_0$  locally uniformly on  $\mathbb{D}$ . In particular,  $h'_\lambda(0) \rightarrow g'_0(0) = 4$ . Since  $h'_\lambda(0)$  is continuous in  $\lambda$  and, by Lemma 11, assumes non-zero real numbers, we also see that  $h'_\lambda(0) > 0$  for all  $\lambda > 0$ . By the uniqueness of the Riemann mapping function and by Lemma 11, we obtain  $g_\lambda = h_\lambda$ . Thus the main theorem has been proved.

## REFERENCES

1. L. V. Ahlfors and L. Bers, *Riemann's mapping theorem for variable metrics*, Ann. of Math. (2) **72** (1960), 385–404.
2. K. Astala and G. J. Martin, *Holomorphic motions*, Papers on analysis, Rep. Univ. Jyväskylä Dep. Math. Stat., vol. 83, Univ. Jyväskylä, 2001, pp. 27–40.
3. Lei Tan (Editor), *The Mandelbrot Set, Theme and Variations*, London Math. Soc. Lecture Note Series, vol. 274, Cambridge Univ. Press, 2000.
4. W. K. Hayman, *Some inequalities in the theory of functions*, Proc. Cambridge Philos. Soc. **44** (1948), 159–178.
5. D. A. Hejhal, *Universal covering maps for variable regions*, Math. Z. **137** (1974), 7–20.
6. E. Hille, *Remarks on a paper by Zeev Nehari*, Bull. Amer. Math. Soc. **55** (1949), 552–553.
7. Y. Imayoshi and M. Taniguchi, *An introduction to Teichmüller spaces*, Springer-Tokyo, 1992.
8. C. I. Kalme, *Remarks on a paper by Lipman Bers*, Ann. of Math. (2) **91** (1970), 601–606.
9. Y. Komori and T. Sugawa, *Bers embedding of the Teichmüller space of a once-punctured torus*, Conform. Geom. Dyn. **8** (2004), 115–142.
10. Y. Komori, T. Sugawa, M. Wada, and Y. Yamashita, *Drawing Bers embeddings of the Teichmüller space of once-punctured tori*, Exper. Math. **15** (2006), 51–60.
11. R. Mañé, P. Sad, and D. Sullivan, *On dynamics of rational maps*, Ann. Sci. École Norm. Sup. **16** (1983), 193–217.
12. Y. Minsky, *The classification of punctured-torus groups*, Ann. of Math. (2) **149** (1999), 559–626.

13. H. Miyachi, *Cusps in complex boundaries of one-dimensional Teichmüller spaces*, Conform. Geom. Dyn. **7** (2003), 103–151.
14. S. Nag, *The Complex Analytic Theory of Teichmüller Spaces*, Wiley, New York, 1988.
15. T. Nakanishi, *A theorem on the outradii of Teichmüller spaces*, J. Math. Soc. Japan **40** (1988), 1–8.
16. ———, *The inner radii of finite-dimensional Teichmüller spaces*, Tôhoku Math. J. (2) **41** (1989), 679–688.
17. T. Nakanishi and J. A. Velling, *A sufficient condition for Teichmüller spaces to have smallest possible inner radii*, Ann. Acad. Sci. Fenn. A. I. Math. **18** (1993), 13–21.
18. T. Nakanishi and H. Yamamoto, *On the outradius of the Teichmüller space*, Comment. Math. Helvetici **64** (1989), 288–299.
19. W. C. Royster, *On the univalence of a certain integral*, Michigan Math. J. **12** (1965), 385–387.
20. H. Sekigawa, *The outradius of the Teichmüller space*, Tôhoku Math. J. (2) **30** (1978), 607–612.
21. T. Sugawa, *The limiting shape of one-dimensional Teichmüller spaces*, in preparation.
22. ———, *Estimates of hyperbolic metric with applications to Teichmüller spaces*, Kyungpook Math. J. **42** (2002), 51–60.

DEPARTMENT OF MATHEMATICS, GRADUATE SCHOOL OF SCIENCE, HIROSHIMA UNIVERSITY, HIGASHI-HIROSHIMA, 739-8526 JAPAN

*E-mail address:* sugawa@math.sci.hiroshima-u.ac.jp